

Best practices: pictorial health warning

Tara Singh Bam

Deputy Regional Director

International Union Against Tuberculosis and Lung Disease (The Union), Singapore

Our future is not written, we have to write our future where everyone can enjoy a healthy and prosperous life

Best Practices Pictorial Health Warnings

First round of 6 pictorial warning labels for factory-made cigarette packs



Second round of 9 pictorial warning labels for factory-made cigarette packs





Pictorial Health Warnings are effective

- Educate people on danger of tobacco use
- Motivate smokers/tobacco users to quit
- Convince youth not to start smoking/tobacco use
- Re-convince ex-tobacco users to remain quitter




Countries requiring picture warnings on cigarette packages-2017





50%+


Countries/jurisdictions requiring warnings of at least 50% of package (on average)



Source: Cigarette packaging health warnings. International Status Report 2016, Canadian Cancer Society Global Tobacco Epidemic 2015. World Health Organization



Pictorial Health Warning in South and South East Asian Countries





Evidences

Larger Pictorial Health Warning



Nepal 90%




Thailand 85%



India 85%



Sri Lanka 80%



Myanmar 75%



Brunei 75%




Lao PDR




Malaysia 55%

Larger pictorial health warnings are effective

PHW made the smokers to reduce cigarettes smoked/day










India Evidences

Remarkable decline of tobacco use




• There is remarkable decline of 54 %, 33 % in the tobacco use among 15-17, 15-24 age groups respectively.


Source: Global Adults Survey 2017



Australia: Evidences proved that PHWs are effective




In Australia, PHW and plain packaging was found effective. Number of adolescents currently smoking reduced by more than 70 per cent, from 23.5% in 1996 to 6.7% in 2014







Myanmar- Comparison between text and pictorial health warning in building public awareness on danger of tobacco use



The Union

Retailers support Pictorial health warnings policies and implementation

Indonesia (N = 1425)


- 84.6% of the retailers support government policy on pictorial health warnings
- 90% of the retailers were found selling cigarettes with pictorial health warning

Nepal (N = 270)

- 92.6% of the retailers support government policy on pictorial health warnings
- 99.3% of the retailers were found selling cigarettes with pictorial health warnings



Indonesia: RI farmers support tougher tobacco control, study says -



Source: http://www.thejakartapost.com/news/2015/10/28/ri-farmers-support-tougher-tobacco-control-study-says.html#sthash.GYBfj6fJ.dpuf

Jakarta Post


Summary

- Larger PHWs are more effective
- All segments of community support PHW; public, youth, retailers and tobacco farmers
- PHW is the most cost effective strategy No cost to government, easy to implement and monitor –
 Government commitments




What do we need to do?

Why not for Indonesia




Indonesia 40%



Nepal 90%



Brunei 75%



Australia Plain Packaging

Cigarettes produced in Indonesia carry a 90% pictorial health warning on packaging for Nepal, 75% PHW for Brunei and plain packaging for Australia

Indonesia: Let's join hands together to increase a size pictorial health warning to at least 75%

40% PHW




Indonesia

90% PHW



Nepal

Plain packaging



Australia

